

## Research Article

# Impact of Harmonic Octave Consonants (Classical Musical Notes) on the Discrete Physiognomic Characters and Different Biochemical Aspects in Sweetleaf viz., *Stevia rebaudiana* (Bertoni)

Sanhita Padhi\*, Rageshree Swain, Pragyan P Rout

Post-Graduation Department of Botany, Ravenshaw University, Cuttack, Odisha

\***Corresponding Author:** Sanhita Padhi, Post-Graduation Department of Botany, Ravenshaw University, Cuttack, Odisha

**Received:** 28 March 2022; **Accepted:** 07 April 2022; **Published:** 06 May 2022

**Citation:** Sanhita Padhi, Rageshree Swain, Pragyan P Rout. Impact of Harmonic Octave Consonants (Classical Musical Notes) on the Discrete Physiognomic Characters and Different Biochemical Aspects in Sweetleaf viz., *Stevia rebaudiana* (Bertoni). International Journal of Plant, Animal and Environmental Sciences 12 (2022): 044-058.

### Abstract

Music is an integral part of our nature and society [1]. The harmonic octave consonants and their frequencies are now-a-days used as a therapy, being popularly called as music therapy. However, the impact of music on our physical and physiological processes of living beings has been acknowledged since ages. Rhythmic and comforting music has an influence on physical and physiological conditions and behavior of living organisms such as humans, plants and animals [2]. According to various studies, Indian Classical music has been confirmed to encourage plant growth

positively and it has also been observed to dominate the other important genres of music such as western classical, rock and monotonous sounds [3]. The present research work is aimed at finding the exclusive impact and effect of the harmonic octave consonants and their frequencies in different strings and closed-pipe Indian classical instrumental music displayed through various *Ragaas*, viz: *Raga Kedar* (flute), *Raga Kedar* (santoor), *Raga-Rageshree* (sitar), *Raga- Bhairavi* (flute), *Raga- Shree* (Sarangi), *Raga- Milan ki Todi* (Sarod), *Raga- Ramkali* (Sitar) on the distinct physiognomic characters and different

biochemical aspects in sweet leaf viz., *Stevia rebaudiana* (Bertoni). The results were surprising as the treated plants showed higher efficacy in terms of both its physiognomic and biochemical aspects as compared to the control plants. The results were depicted here through tabulations and figures.

Physiognomic parameters like plant height, number of leaves per plant, leaf lamina length, leaf lamina breadth, leaf texture, leaf color, spread of plant in east- west and in north-south directions and diameter of the stem for a period of 30 days. Treated plants showed an increment in the growth as compared to the control and survived longer (20 days longer) than the control plants.

For instance, in sweet leaf, the average no. of leaves in control was 53 per plant and 64 in treated plants. Similarly, the average height of the treated (experimental) plant was found to be greater i.e., 28 cm as compared to the lower average height of 21 cm as shown by the control (untreated) plant samples. The biochemical analysis also revealed some promising observations. On performing the biochemical experiments with the leaves of sweet leaf (both control and experimental), the total protein, carbohydrate and chlorophyll contents in the treated leaves was found to be much higher i.e., 142.25 mg/gm, 114 mg/gm and 34.69 µg/gm fresh weight, respectively. The control plants showed much lesser values such as 87.5 mg/gm, 79 mg/gm and 16.44 µg/gm for total protein, total carbohydrate and total chlorophyll content respectively. The leaves of treated (experimental) sweet leaf showed a increased value of 75.66 mg/100gm whereas the control leaves showed 42.52 mg/100 gm of ascorbic acid.

**Keywords:** Sound; Music; Indian Classical *Ragas*; Harmonic octave consonants; Strings and closed-pipe instruments; Physiognomic study; Biochemical study

## Abbreviations

Hz=Hertz; I=Sound Intensity; L=Sound Intensity Level; Db=Decibel

## 1. Introduction

### 1.1 Sound

Sound is a form of energy. It is a compressional wave triggered by vibrations [4]. The vibrations that are felt on touching the surface of a speaker are equal to the vibrations disturbing the air. Sound is similar to countless other things which are invisible in our universe. It cannot be seen as well but sound can be heard as it reaches our ears in the form of waves produced when particles in the medium vibrates [5]. Hence, vibration in a body generates sound.

### 1.2 Music

Music and noise are two different things; the former has a soothing effect and the latter is infuriating. Music is related to rhythmic and periodic vibrations while noise has no periodic vibrations [1]. Music for about a thousand of years has been a source of harmony for people. It is a soothing and tranquilizing form that is produced from musical instrument. Music is a fine art of sound and is basically prearranged by people to express their feelings a certain way.

Music certainly has many dimensions in stirring various physical, psychological, spiritual and social stages of awareness [6]. Currently, music is categorized as both positive and negative. Music that

has valuable potentials and inspires at emotional and spiritual levels and causes relaxing, calming and healing effects is known as positive music whereas negative music induces negative emotions, irritation, bitterness, sadness, animosity and terror [2].

### 1.3 Octave

Octave which can simply be known as a doubling in frequency (40 Hz is one octave greater than 20 Hz) is one dynamic conception related to music. According to Shah (2001), the several 'octaves' has been arranged on the basis of audible sound range and depending on some mathematical principles, an individual octave has been classified into intervals. An octave is categorized into twelve parts, which is the width of a semitone, i.e. the frequency ratio of the interval between two adjacent notes, is the twelfth root of two and is mathematically expressed as [7]:

$$12\sqrt[12]{2} = 2^{1/2} \approx 1.059463, \text{ which is again equivalent to } e^{1/2 \ln 2} \approx 1.059463$$

This interval is divided into 100 cents. An equivalent of octave in Indian Classical Music is Saptak that is a constituent of twenty-two tones stated to as '*Shruti*'. The seven pure notes in Indian Classical Music are called '*shuddha swaras*' and are composed by the microtones i.e. *shrutis* showing distinct frequency range (Table 1).

After the first concept of perception of sound waves by plants as suggested by Sir Jagdish Chandra Bose, a Nobel Prize winner in 1927, who was renowned for his work on the physiology of plants; Dorothy Retallack also carried out one of the first experiments in 1973 to study the relationship between plants and

music. Various styles of music were used by Retallack in her experiments and she discovered that the plants showed a tendency to move away from Led Zeppelin and Jimi Hendrix but Bach organ music and jazz attracted them. Nevertheless, she found that North Indian classical music played on the sitar was their favorite but country music was seen to have no such relevance in plant life [3].

### 1.4 Music in relation to plants

All living beings, be it animals or plants, all respond to external stimuli. Plants, the multicellular organisms also respond to various types of external stimuli including Sound vibration. These vibrations stimulate different phytochemical and biochemical reactions which in turn helps in the developmental processes of the plants. In case of human beings, sound waves set up vibration in our ear drum which in turn is perceived by the brain and help us to recognize the type of sound of varying frequencies and amplitude.

Plants, in a similar way receive vibrations through protoplast. Various studies related to the effects of sound waves subjecting to seeds and plants, generally known as sonication, has been published in scientific literature [8, 9]. The effect of music on 30 Rose (*Rosa chinensis*) plants taken in separate pots was studied [10].

The plants exposed to Indian classical music exhibited the highest internode elongation, which evidently displayed that exposing the plants to Vedic chants and Indian classical music stimulates the growth of plants as compared to the control group and the plants those were subjected to Western pop and Rock music.

### 1.5 Musical notes and shruti

A total of 240 vibrations per second are produced by the note Shadja (Sa). Likewise, number of vibrations generated by other notes is: Rishab (Re)-270,

Gandhar (Ga)-300, Madhyam (Ma)-320, Pancham (Pa)-360, Dhaivat (Dha)-400 and Nishad (Ni)-450 [11].

Shruti	Frequency ratio ( $f = v/\lambda$ )	Frequency (Hertz)
Sa	1/1	240
Re1	32/31	252.8
Re2	16/15	256
Re3	10/9	266.6
Re4	9/8	270
Ga1	32/27	284.4
Ga2	6/5	288
Ga3	5/4	300
Ga4	81/64	303.7
Ma1	4/3	320
Ma2	27/20	324
Ma3	45/32	337.5
Ma4	64/45	341.3
Pa	3/2	360
Dha1	128/81	379
Dha2	8/5	384
Dha3	5/3	400
Dha4	27/16	405
Ni1	16/9	426.6
Ni2	9/5	432
Ni3	15/8	450
Ni4	31/16	465

Sa - Shadja, Re - Rishab, Ga - Gandhar, Ma- Madhyam, Pa – Pancham, Dha- Dhaivat and Ni - Nishad (Ni), f- frequency ratio, v- speed,  $\lambda$ -lambda

**Table 1:** Frequency ratios of ‘shruti’ values (Source: Shah, 2001).

## 2. Materials and Methods

### 2.1 Materials

**2.1.1 Acoustic chamber:** An acoustic chamber was used with a sound system to produce the sound of known frequency through Indian Classical Instrumental Music both String music and Closed pipe music) for the experimental plants.

**2.1.2 Plant material:** The plant commonly called as sweet leaf [12] with the scientific name *Stevia rebaudiana* (Bertoni) was purchased from medicinal garden at Patrapada, Bhubaneswar in large number and used as for the experiments.

### 2.1.3 Measuring tape and thread

**2.1.4 Ragaas:** The application of the Indian Classical Instrumental music notes (Octaves) through different *Ragaas* were applied to our experimental plant set in the following time sequence: -

- Raga Kedar- Flute (closed pipe), by Pandit Pannalal Ghosh (38:14) during early morning between 6am-8am, followed by-
- Raga Kedar- Santoor (stringed instrument), by Pandit Shiv Kumar Sharma (23:14) in the Morning time between 10am-12pm,
- Raga Rageshree- Sitar (stringed instrument), by Nikhil Banerjee (29:15) in the evening between 4pm-6pm followed by-
- Raga Bhairavi- Flute (closed pipe), by Pandit Hari Prasad Chaurasia (9:50)

All the above mentioned four *Ragaas* have been observed to promote seed germination and vegetative

growth. Physiognomic studies were conducted on an interval of four days for a period of 30 days. A thread was used to measure the height of the plant, leaf lamina length, leaf lamina breadth, and spread of the plant in east-west and north-south directions. It was then placed over a 60 cm long measuring tape and the measurement data were recorded in table.

### 2.2 Methodology

The plants were treated with the above harmonic sound frequencies of musical notes for 4 hours at frequent intervals daily.

**2.1.1 Physiognomic study:** All the plants were kept inside the acoustic chamber. The control plants were taken outside of the chamber when the treatment was given. Different aspects of physiognomic studies like height of the plant, leaf lamina length per plant, leaf lamina breadth per plant, spread of the plant in east-west direction and in north- south direction, and diameter of the stem were considered. These aspects were measured in every four days.

**2.1.2 Biochemical analysis of leaves:** To study the effect of this Indian Classical instrumental Music on biochemical aspects through various observations and inferences from the experiments, a series of biochemical tests were performed in order to estimate the quantity of total proteins, total carbohydrates and total chlorophyll contents in the leaves of *Stevia*, treated with *ragaas* verses control.

**2.1.2.1 Protein estimation:** The amount of protein was quantified by using the method of Lowry *et al* 1951 [13]. 500mg per 100gm fresh weight of both control and treated leaf sample was taken for analysis.

**2.1.2.2 Carbohydrate estimation:** Anthrone method or Hedge method, 1962 [14] was used for carbohydrate estimation. 100mg per 100gm fresh weight of both control and treated leaf sample was taken for analysis.

**2.1.2.3 Chlorophyll estimation:** Chlorophyll is extracted in 80% acetone and the absorbance are read at 663 and 645 nm in a spectrophotometer, using the absorption co-efficient, the amount of chlorophyll is calculated [15]. 500mg per 100gm fresh weight of leaf sample was taken for analysis.

Formulas

The amount of chlorophyll present was calculated by the following formulae,

$$\text{Mg chl a /gm fresh weight} = 12.7(A663) - 2.69(A645)*V/1000*W$$

$$\text{Mg chl b / gm fresh weight} = 22.9(A645) + 4.68(A663)*V/1000*W$$

$$\text{Mg total chl/gm fresh weight} = 20.2(A645) + 8.02(A663)*V/1000*W$$

#### 2.1.2.4 Estimation of ascorbic acid:

Ascorbic acid is estimated by DCPIP method (Harris and Ray, 1935) in both control and treated leaf sample. 500mg per 100gm fresh weight of both control and treated leaf sample of sweet leaf was taken for analysis.

Ascorbic acid content (mg/100g) =  $0.5 * V_2 / V_1 * \text{total volume of stock solution} / \text{vol. taken for analysis} * 100 / \text{wt. of sample taken (g)}$ , Where,  $V_1$  = volume of dye consumed for standard (ml),  $V_2$  = volume of dye consumed for sample (ml).

### 3. Results

Several studies have proved that there is an association between sound vibrations and plants lives and that both are profoundly interconnected to each other [3]. While sound with low amplitude improves plant growth, likewise, any type of sound vibrations with large amplitude values for e.g., rock music declines plant growth [3]. The present research work embraces two important aspects of the plant growth and the physiological processes involved therein were taken into consideration:

- Effect of harmonic octave consonants on the physiognomy of sweet leaf, *Stevia rebaudiana* (Bertoni).
- Effect of harmonic octave consonants on the biochemical analysis of sweet leaf, *Stevia rebaudiana* (Bertoni).

#### 3.1 Effect on physiognomy of plants

Physiognomy is amalgam of external features and different growth forms in plants. It is one of the most pivotal studies in case of plant classification.

##### 3.1.1 Physiognomic study of *Stevia rebaudiana*

**(Bertoni):** To perform the experiment two sets were taken; one set with 4 plants as control and the other set with 4 plants as treated. Physiognomic parameters like plant height, no. of leaves per plant, leaf lamina length, leaf lamina breadth, leaf texture, leaf color, spread of plant in east- west and in north-south directions and diameter of the stem for a period of 30 days. Treated plants showed an increment in the growth as compared to the control and survived longer (20 days longer) than the control plants. The


average no. of leaves in control was 53 per plant and 64 in treated plants and the rest all data are given in tabular form. The observations are shown in tables

and figures (Table 2; Figure 1; Figure 2; Figure 3; Figure 4).

Physiognomic parameters	Control (average) (cm)	Treated (average) (cm)
Ph	21.5	28
Sop-(E-W)	22.67	25.71
Sop-(N-S)	21.5	28
Lll	2.34	3.5
Llb	1.2	1.3
Dos	1.8	1.9

Ph-Plant height, Sop-(E-W)- Spread of plant (East-West), Sop-(N-S)- Spread of plant (North- South), Lll- Leaf lamina length, Llb- Leaf lamina breadth, Dos- Diameter of Stem

**Table 2:** Measurements of physiognomic parameters in sweet leaf (*Stevia rebaudiana*) after 30 days of treatment with music.




**Figure 1:** Measurements of physiognomic parameters in sweet leaf (*Stevia rebaudiana*).





**Figure 2:** Sweet leaf (*Stevia rebaudiana*) plants before treatment (A1-A4).



**Figure 3:** Treated sweet leaf (*Stevia rebaudiana*) set up after the period of 30 days (B1-B4).





**Figure 4:** Control sweet leaf (*Stevia rebaudiana*) plants after 30 days (C1-C4).

**3.2 Effects of harmonic octave consonants on the biochemical aspects of sweet leaf (*Stevia rebaudiana*)**


**3.2.1 Protein estimation**

**3.2.1.1 Effect of harmonic octave consonants on total protein contents:** Leaves of both control and

treated were taken for extract preparation. The total protein content of the leaf samples was estimated by the Lowry et al 1951 [13] method. The data generated is recorded in table (Table 3) and analyzed in figure (Figure 5).

Treatment	Total protein content (500mg/100gm fresh weight)
Control	87.5
Treated	142.25

**Table 2:** Total protein content of control and treated plants of sweet leaf (*Stevia rebaudiana*).



**Figure 5:** Total protein content of control and treated plants of sweet leaf (*Stevia rebaudiana*).

### 3.2.2 Total carbohydrate estimation


#### 3.2.2.1 Effect of harmonic octave consonants on total carbohydrate content:

Young leaves were selected from the treated plants. The extract was prepared from leaves of both control and treated plants and the total carbohydrate was estimated using the Anthrone method or Hedge method,

1962 [14]. Equal amount of leaf sample was taken for both control and experiment. The carbohydrate concentration was found higher in the treated plants as compared to the control plants. The data are recorded in table (Table 4) and analyzed in figure (Figure 6).

Treatment	Total carbohydrate content (100mg/100gm fresh weight)
Control	79
Treated	114

**Table 4:** Total carbohydrate content of control and treated plants of sweet leaf (*Stevia rebaudiana*).



**Figure 6:** Total carbohydrate content of control and treated plants of *Stevia rebaudiana*.

### 3.2.3 Chlorophyll estimation

**3.2.3.1 Effect of harmonic octave consonants on chlorophyll content:** The control and treated plants were subjected to chlorophyll estimation. The acetone method was used for estimation of chlorophyll content. For calculations of chlorophyll a, chlorophyll b and total chlorophyll

content Arnon’s (1949) formulae were used. The chlorophyll a chlorophyll b and total chlorophyll content were found effectively higher in treated plants than in control plants. The data are recorded in table (Table 5) and analyzed through figure (Figure 7).

Treatment	Chlorophyll a (500µg/100gm fresh weight)	Chlorophyll b (500µg/100gm fresh weight)	Total chlorophyll content (500µg/100gm fresh weight)
Control	17.75	19.70	16.44
Treated	24.86	38.78	34.69

**Table 5:** Chlorophyll a, Chlorophyll b, and total chlorophyll content of sweet leaf (*Stevia rebaudiana*).




Figure 7: Chlorophyll a, Chlorophyll b, and total chlorophyll content of sweet leaf (*Stevia rebaudiana*).


**3.2.4 Estimation of ascorbic acid in *Stevia rebaudiana* by DCPIP method:**

Ascorbic acid, also known as vitamin C is necessary for wound healing and many other functions in the body. Sweet leaf also shows some amount of ascorbic acid i.e., 14.98 mg·100 g<sup>-1</sup> [16]. Because of its immense importance and also due to its high presence, this analysis was

carried out in the plant, *Stevia rebaudiana*. The analysis was carried out in the leaf of *Stevia rebaudiana*. Leaf of the treated plant showed an elevated quantity of the ascorbic acid as compared to that in the control plants. The data has been presented in the table (Table 6) and figure (Figure 8) below.

Plant parts	Control	Treated
Leaf	42.52	75.66

Table 6: Ascorbic acid content (500mg/100gm) in sweet leaf (*Stevia rebaudiana*).



**Figure 8:** Ascorbic acid content (500mg/100gm) in control and treated leaf sample of sweet leaf (*Stevia rebaudiana*).

#### 4. Discussion

Audible sound (20–20000Hz) widely exists in natural world. However, the interaction between audible sound and the growth of plants is usually scanty in biophysics research. Not much effort has been put forth in studying the relation of plant processes and audible sound.

A certain study has revealed that plants have mechano-sensitive channels that perceive sound vibrations. In another experiment it was shown that sound vibrations can alter tensions in biological membranes that could possibly stimulate signaling pathways through the activation of the mechano-sensitive channels in the membranes of plant cell (Haswell and Meyerowitz, 2006). Other experiments carried out showed that certain frequencies of sound vibrations can induce seed germination, root

elongation, callus growth, and cell [17]. An experiment was conducted in which 12 tobacco plants were exposed to noise and showed overall 40.6 % decrease as compared to the control plants [18].

In the present research study, it was also seen that there was an increment in the overall growth of the plant body compared to the control though not much of difference was found. *Stevia rebaudiana* (Bertoni) when subjected to soft harmonic sound vibrations, the results showed a difference i.e., there was an increment in the physiognomic parameters of the treated plant as compared to the control (untreated) plant. The possible explanation to this would be the perception of sound vibration and relay of extended signaling pathways leading to the overall increased growth of the plant and leaves.

Audible harmonic frequencies not only accelerate growth but also significantly influences the concentration of various metabolites; e.g., chlorophyll and starch are increased by it [19]. Also, musical vibration stimulates water molecules within biological systems [20]. As a result, the temperature, which is a measure of motion of molecules is raised due to vibration of the water molecules. The rate of metabolism is thereby, increased with the increase in temperature. Resonances from the audible harmonic sound vibration can combine directly into biological systems because they are constituted mostly of water.

Indole Acetic Acid (IAA) is an essential plant hormone that helps in plant's growth and development. Zhu and co-workers observed that IAA content in plants were found at an increased level in six species of vegetable plants when exposed to musical acoustic frequencies in comparison to the control plants [21]. The present study also revealed an increase in carbohydrates, proteins and chlorophyll content. The increased metabolic rate might be a reason for the observed results.

### 5. Acknowledgements

The authors would like to sincerely acknowledge the Department of Botany, Ravenshaw University, for providing all the supports and the infrastructural facilities required to carry out the entire work. We also acknowledge the Acoustical and Biochemistry laboratory where the entire experimental work was done. Our special thanks and acknowledgement go to our supervisor Prof. Sanhita Padhi for her immense support and timely supervision.

### References

1. Lapp DR. The physics of music and musical instruments. Wright Center for Innovative Science Education. Tufts University, Medford, Massachusetts (2002): 119.
2. Ekici N, Dane F, Mamedorah L, et al. The effects of different musical elements on root growth and mitosis in onion (*Allium cepa*) root apical meristem (musical and biological experimental study). *Asian J. Plant Sci* 6 (2007): 369-373.
3. Retallack D. *The Sound of Music and Plants*. Santa Monica, CA: DeVorss and Co. (1973): 96.
4. Wood A. *The physics of music*. Chapman and Hall, London (1976): 258.
5. Rigden JS. *Physics and the sound of music*. Wiley, New York (1977): 71.
6. Kneafsey R. The therapeutic use of music in a care of the elderly setting: a literature review. *J. Clin. Nurs* 6 (1997): 341-346.
7. Kuttner FA. Prince Chu Tsai-Yü's Life and Work: A Re-Evaluation of His Contribution to Equal Temperament Theory. *Ethnomusicology* 19 (1975): 163-113.
8. Suslick KS. *Ultrasound: Its chemical, physical and biological effects*. New York, VCH Publishers (1989): 227.
9. Joersbo M, Brunstedt J. Sonication a new method for gene transfer to plants. *Physiol. Plant* 85 (1992): 230-234.
10. Chivukula V, Ramaswamy S. Effect of different types of music on *Rosa chinensis* plants. *Int. J. of Environ Sci Te* 5 (2014): 1-4.

11. Thakkar MG, Kajal C. Science of Ragas: The control on living beings to the cosmic elements. *Journal of Environmental Research and Development* 9 (2014): 260-270.
12. Marcinek K, Krejpcio Z. Stevia rebaudiana Bertoni-chemical composition and functional properties. *Acta Sci. Pol. Technol. Aliment* 14 (2015): 145-152.
13. Tambe SS, Deore S, Ahire PP, et al. Biochemical evaluation of some medicinal plant of Marathwada region in Maharashtra, *International Journal of Pharmaceutical Research and Bio-Science* 1 (2012): 185-194.
14. Das BK, Choudhury BK, Kar M. Quantitative estimation of changes in Biochemical constituents of Mahua (*Madhuca indica* syn. *Bassia latifolia*) flowers during postharvest storage. *Journal of Food processing and preservation* 34 (2010): 831-844.
15. Arnon DI. The estimation of chlorophyll in plants. *Journal of Plant Physiology* 24 (1949): 1-15.
16. Bugaj B, Leszczyńska T, Pysz M, et al. Charakterystyka i właściwości prozdrowotne *Stevia rebaudiana* (Bertoni) [Characteristics and health-promoting properties of *Stevia rebaudiana* (Bertoni)]. *Żywn. Nauka Techn. Jakość* 3 (2013): 27-38.
17. Gagliano M. Green symphonies: a call for studies on acoustic communication in plants. *International Society for Behavioral Ecology* 24 (2012): 789-796.
18. Woodlief CB, Royster LH, Huang BK. Effect of random noise on plant growth. *The Journal of the Acoustical Society of America* 46 (1969): 481-482.
19. Sharma D, Gupta U, Fernandes AJ, et al. The effect of music on physico-chemical parameters of selected plants. *Int. J. of Plant, Animal and Environmental Sciences* 5 (2015): 282-287.
20. Creath K, Schwartz GE. Measuring effects of music, noise, and healing energy using a seed germination bioassay. *J Altern Complem Med* 10 (2004): 113-122.
21. Jun-ruZ, Shi-renJ, Lian-qing S. Effects of music acoustic frequency on Indoleacetic Acid in plants. *Agricultural Science and Technology* 12 (2011): 1749-1752.



This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution \(CC-BY\) license 4.0](https://creativecommons.org/licenses/by/4.0/)